

No aspirin or salicylic acid in children of any age

Date:

Age:

Weight:

- Keep all medicines out of reach of children
- No Acetaminophen for children less than 3 months old
- For fever more than 100 degrees in children less than 3 months old, call the doctor
- Acetaminophen and Tylenol® are the same
- Give the dose for your child's weight as listed below
- Give every 4 hours for fever more than 100.4 degrees
- Other medicines may have Acetaminophen in them. Do not give more Acetaminophen than ordered

Weight (pounds)	Infant Concentrated Drops 160 mg per 5 ml	Children's 160 mg per 5 ml	Chewable Tablets 160 mg per tablet
6 to 23	Call doctor		
24 to 35	5 ml	5 ml (1 tsp)	1 tablet
36 to 47		7.5 ml	1½ tablet
48 to 59		10 ml (2 tsp)	2 tablets
60 to 71		12.5 ml	2½ tablets
72 to 95		15 ml (3 tsp)	3 tablets
more than 95		20 ml (4 tsp)	4 tablets

Give your child the right dose

- Be sure to read the right dose for your child
- **Only** use the device you get with the medicine to give the right dose

tsp = teaspoon

ml = milliliter

1 tsp = 5 ml



Dosing information for Acetaminophen and Ibuprofen from Tylenol for Health Professionals <https://www.tylenolprofessional.com/dosage.html>

Carolinas HealthCare System complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability or sex.

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1- 800-821-1535.

CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1- 800-821-1535.

